

Antimicrobial Activity of Marine Samples Collected From The Different Coasts of Turkey

Belma KONUKLUGİL^{1,*}, Bülent GÖZCELİOĞLU²

¹Ankara University, Faculty of Pharmacy, Department of Pharmacognosy, Döğol Cad., 06100 Ankara, TURKEY, ²The Scientific and Technological Research Council of Turkey (TÜBİTAK), Kavaklıdere, 06100 Ankara, TURKEY

During our search for bioactive compounds from Turkish marine sources, we have detected antimicrobial activity of some of the extracts that have been prepared from sponges, and tunicate and soft corals from different coasts of Turkey. All samples were extracted with methanol. Antimicrobial activity test was conducted against 5 microbial pathogens; (two gram positive bacteria, one yeast and two gram negative bacteria) methicillin-resistant *Staphylococcus aureus* (ATCC 33591), vancomycin-resistant *Enterococcus faecium* (EF379), *Candida albicans* (ATCC 14035), *Pseudomonas aeruginosa* (ATCC 14210), *Proteus vulgaris* (ATCC 12454), respectively. While crude extracts and filtrated *Axinella verrucosa* extract showed strong antimicrobial activity, some of the sponge species showed strong or modest activity depending on their localities and concentrations.

Key words: Marine sponge, Tunicate, Soft coral, Secondary metabolite, Antimicrobial activity

Türkiye'nin Farklı Kıyılarından Toplanan Denizel Örneklerin Antimikrobiyal Aktiviteleri

Türkiye denizlerinde bulunan süngerlerdeki bioaktif maddelerle ilgili devam eden çalışmalarımız kapsamında, bu çalışmada değişik kıyılardan toplanan sünger, tunikat ve yumuşak mercan örneklerinin metanollü ekstrelerinin antimikrobiyal etkisi incelenmiştir. Antimikrobiyal çalışma için 5 mikrobiyal patojen iki Gram (+) bakteri, iki Gram (-) bakteri ve bir mantar metisillin dirençli-*Staphylococcus aureus* (ATCC 33591), vankomycin- dirençli *Enterococcus faecium* (EF379), *Candida albicans* (ATCC 14035), *Pseudomonas aeruginosa* (ATCC 14210) ve *Proteus vulgaris* (ATCC 12454) kullanılmıştır. Konsantrasyon ve lokaliteye bağlı olarak sünger örneklerinin kuvvetli veya orta duyarlı antimikrobiyal etki gösterdiği tespit edilmiştir. İncelenen türler içinde *Axinella verrucosa*'nın hem ham ekstresinde hemde süzölmüş ekstresinde güçlü antimikrobiyal aktivite gösterdiği bulunmuştur.

Anahtar kelimeler: Deniz süngeri, Tunikat, Yumuşak mercan, Sekonder metabolitler, Antimikrobiyal etki

*Correspondence: E-mail: belma.konuklugil@gmail.com; Tel: +903122033092

INTRODUCTION

The ocean which covers almost 70 % of the planet's surface provide a huge biodiversity with potential as immeasurable source of natural products. Until now more than 17000 marine secondary metabolites have been described of which sponges are responsible for more than 5300 different products (1). The chemical diversity of sponge metabolites is notable, in addition to the unusual nucleosides, terpenes, sterols, peptides, alkaloids, fatty acids and etc. (2). Several marine natural products have a significant biological activity and many of them are currently, in different phases of clinical trials as drug candidates. In the United States there are three FDA approved marine derived drugs, namely cytarabine, vidarabine and ziconotide. Currently, trabectedin has been approved by the European Agency for Evaluation of medicinal products (EMA) and is completing key Phase III studies in US for approval (3). Marine sponges are among the richest sources of pharmacologically active compounds from marine organisms. Many interesting secondary metabolites have been isolated from various species of sponges including powerful antiviral, antimalarial, antitumor and anti-inflammatory agents (4,5). Screening of marine sponge extracts for antibacterial activity led to the isolation and characterization of a wide range of active compounds, (alkylpiperidine, bromopyrrole and pyrroloiminoquinone alkaloids, sesquiterpene-quinones /- hydroquinones, terpenoids, phenolic compounds, peptides and proteins, polyketide and polysaccharides) including some promising therapeutic leads (6,7). The extracts of different sponge species were also found to be active against a wide spectrum of bacterial strains isolated from hospitalized human patients (8). Extensive research has been done to unveil the antimicrobial compounds of sponges including different genera such as, *Axinella*, *Ircinia*, *Agelas*, *Dysidea* and results are amazingly diverse and productive (9-13).

Screening of organic extracts from marine sponges is a common approach to identify biomedically important compounds. The objective of our study was screening of

methanolic extracts of 33 marine samples collected from different coasts of Turkey for antimicrobial activities with the aim of identifying novel compounds with interesting and potentially useful therapeutic activities.

EXPERIMENTAL

Marine Samples

Sponge species (*Ircinia* sp., *Petrocia ficiformis*, *Dysidea avara*, *Agelas oroides*, *Axinella verrucosa*, *Aplysina aerophoba*, *Chondrilla nucula*, *Agelas oroides*, *Axinella polypoides*, *Axinella damicornis*, *Sarcotragus spinulosa*, *Cicalypta carballoi*, *Ircinia fasciculata*, *Chondrosia reniformis*), soft coral (*Eucinella cingularis*) and tunicate (*Aplidium elegans*) were collected by scuba divers in different coasts of Turkey (Fethiye, Ayyalık, Danaadası, Güvercinlik, İbrice, Kaş, Kemer, Sinekli, Turgutreis) in March 2012, and were identified by Dr. Bülent Gözcelioğlu (one of the authors). Sponge samples, soft coral, and tunicate were deposited at Ankara University, Faculty of Pharmacy, Ankara, Turkey.

Preparation of the extracts

Fresh sponge samples were chopped into small pieces and extracted individually with methanol (3 x 50 mL) for several times at room temperature. The extracts were filtered and evaporated in vacuo until dryness. Methanolic extracts from 33 marine samples were evaluated for their antimicrobial properties. In order to obtain LC/MS data on these samples, an aliquot of each extract was filtrated on C18 Sep-Pack cartridge using MeOH as eluent. Microbroth dilution assays were carried at a concentration of 250 µg/mL using both original crude extracts and C18 fractions against 5 microbial pathogens: two gram positive bacteria methicillin-resistant *Staphylococcus aureus* (MRSA) and vancomycin-resistant *Enterococcus* (VRE), one yeast *Candida albicans* fermented in non-shaking (CA) and shaking conditions (CA_S) and, two Gram negative bacteria *Pseudomonas aeruginosa* (PA) and *Proteus vulgaris* (PV) (14).

RESULTS AND DISCUSSION

Strong antimicrobial activities were observed against gram positive bacteria for a few number of crude extracts at a concentration of 250 µg/mL. Moreover, MeOH filtration on C18 Sep–Pack improved the amount of secondary metabolites in the fractions by removing salts and lipid-like molecules and resulted in an increase of the hits rate. Sample species, their localities and % inhibition values of crude extracts and filtrated extracts are shown in Tables 1, 2, 3 by the concentrations of 250 µg/mL and 50

µg/mL, respectively.

Tables show that crude extract of *Axinella polypoides* has strong antimicrobial activity against MRSA (101%), VRE (96%), CA (101%), PA (99%) and PV (100%). *Axinella verrucosa* and *Dictyonella incisa* collected from Fethiye, showed strong antimicrobial activity against MRSA (101% *Axinella verrucosa*, 81% *Dictyonella incisa*); *Axinella verrucosa* showed strong activity against VRE (100%), as well. On the other hand, *Cicalypta carballoi* collected from Kaş showed moderate activity against VRE (54%).

Table 1. Species, localities and % inhibition values of extracts 5X (250 µg/mL)

[Methicillin-resistant *Staphylococcus aureus* (MRSA), Vancomycin-resistant *Enterococcus faecium* (VRE), *Candida albicans* (CA), *Pseudomonas aeruginosa* (PA), *Proteus vulgaris* (PV)]

	Taxonomy	Location	MRSA	VRE	CA	CA_S	PA	PV
1	<i>Ircinia oros</i>	Fethiye 1	0	0	7	0	7	0
2	<i>Axinella verrucosa</i>	Fethiye 5	101	100	27	23	26	3
3	<i>Petrocia ficiformis</i>	Ayvalık 1	0	0	0	0	0	13
4	<i>Ircinia oros</i>	Ayvalık 2	0	0	0	0	0	16
5	<i>Dysidea avara</i>	Ayvalık 3	0	0	6	0	6	1
6	<i>Agelas oroides</i>	Ayvalık 4	0	0	0	11	0	4
7	<i>Axinella verrucosa</i>	Ayvalık 5	0	0	0	15	0	10
8	<i>Aplysina aerophoba</i>	Ayvalık 6	0	0	1	0	0	0
9	<i>Chondrilla nucula</i>	Ayvalık 10	0	0	3	0	2	0
10	<i>Agelas oroides</i>	Ayvalık 13	0	0	10	0	10	10
11	<i>Axinella polypoides</i>	Danaadası	0	0	13	4	13	0
12	<i>Petrocia ficiformis</i>	Danaadası	0	0	15	30	15	15
13	<i>Ircinia oros</i>	Fethiye 1	0	0	13	17	13	14
14	<i>Petrocia ficiformis</i>	Fethiye 2	19	0	0	30	1	19
15	<i>Agelas oroides</i>	Fethiye 3	0	0	18	6	17	8
16	<i>Axinella damicornis</i>	Fethiye 4	0	3	13	32	12	9
17	<i>Axinella verrucosa</i>	Fethiye 5	13	0	14	0	13	0
18	<i>Dysidea avara</i>	Fethiye 7	10	12	15	0	14	1
19	<i>Sarcotragus spinulosa</i>	Fethiye 10	3	0	0	0	0	0
20	<i>Axinella polypoides</i>	Fethiye 11	100	96	101	0	99	100
21	<i>Ircinia variabilis</i>	Fethiye 13	0	0	9	0	9	8
22	<i>Petrocia ficiformis</i>	Fethiye 14	0	0	12	0	11	9
23	<i>Petrocia ficiformis</i>	Fethiye 14	0	0	0	0	0	17
24	<i>Dictyonella incisa</i>	Fethiye 15	81	0	15	1	15	12
25	<i>Chondrilla nucula</i>	Güvercinlik	0	0	19	23	18	0
26	<i>Dysidea avara</i>	İbrice 4	0	41	21	23	20	0
27	<i>Agelas oroides</i>	Kaş 2	0	0	17	0	17	0
28	<i>Cicalypta carballoi</i>	Kaş 5	0	54	8	0	7	8
29	<i>Ircinia fasciculata</i>	Kemer 3	0	8	0	0	0	0
30	<i>Cliona viridis</i>	Sinekli 8	0	0	0	0	0	14
31	<i>Ircinia variabilis</i>	Sinekli 9	3	0	9	4	9	15
32	<i>Agelas oroides</i>	Turgutreis 17	0	0	16	15	16	10
33	<i>Chondrosia reniformis</i>	Turgutreis 23	0	0	17	10	16	41

Table 2. Species, localities and % inhibition values of filtrated extracts 5X (250 µg/mL)

ID	Taxonomy	Location	MRSA	VRE	CA	CA_S	PA*	PV*
1	<i>Ircinia oros</i>	Fethiye 1	3	0	0	-	3	7
2	<i>Axinella verrucosa</i>	Fethiye 5	6	2	0	-	0	20
3	<i>Petrocia ficiformis</i>	Ayvalık 1	0	0	0	-	0	9
4	<i>Ircinia oros</i>	Ayvalık 2	11	1	0	-	0	15
5	<i>Dysidea avara</i>	Ayvalık 3	0	70	0	-	1	11
6	<i>Agelas oroides</i>	Ayvalık 4	13	0	0	-	17	13
7	<i>Axinella verrucosa</i>	Ayvalık 5	0	0	0	-	9	0
8	<i>Aplysina aerophoba</i>	Ayvalık 6	2	0	0	-	17	0
9	<i>Chondrilla nucula</i>	Ayvalık 10	16	0	0	-	0	0
10	<i>Aplidium elegans</i>	Ayvalık 11	0	0	0	-	0	0
11	<i>Eucinella singularis</i>	Ayvalık 12	0	0	0	-	5	0
12	<i>Agelas oroides</i>	Ayvalık 13	0	0	0	-	0	20
13	<i>Axinella polypoides</i>	Danaadası (32m)	10	0	0	-	10	0
14	<i>Petrocia ficiformis</i>	Danaadası	0	0	0	-	16	0
15	<i>Ircinia oros</i>	Fethiye 1	43	0	0	-	9	0
16	<i>Petrocia ficiformis</i>	Fethiye 2	0	0	0	-	7	0
17	<i>Agelas oroides</i>	Fethiye 3	98	96	0	-	12	95
18	<i>Axinella damicornis</i>	Fethiye 4	0	0	0	-	5	0
19	<i>Axinella verrucosa</i>	Fethiye 5	100	100	0	-	5	0
20	<i>Dysidea avara</i>	Fethiye 7	94	87	0	-	28	0
21	<i>Sarcotragus spinulosa</i>	Fethiye 10	61	0	0	-	0	0
22	<i>Axinella polypoides</i>	Fethiye 11	18	0	0	-	22	47
23	<i>Ircinia variabilis</i>	Fethiye 13	10	4	0	-	0	4
24	<i>Petrocia ficiformis</i>	Fethiye 14	14	10	0	-	19	9
25	<i>Petrocia ficiformis</i>	Fethiye 14	0	0	0	-	0	8
26	<i>Dictyonella incisa</i>	Fethiye 15	100	95	0	-	7	0
27	<i>Chondrilla nucula</i>	Güvercinlik	3	0	0	-	15	0
28	<i>Dysidea avara</i>	İbrice 4	82	62	0	-	3	0
29	<i>Agelas oroides</i>	Kaş 2	66	91	0	-	0	0
30	<i>Ciocalypa carbolloi</i>	Kaş 5	41	75	0	-	30	0
31	<i>Ircinia fasciculata</i>	Kemer 3	77	26	0	-	0	0
32	<i>Agelas oroides</i>	Turgutreis 17	17	0	0	-	20	0
33	<i>Chondrosia reniformis</i>	Turgutreis 23	0	0	0	-	0	51

Filtrated extracts (250 µg/mL) of *Dysidea avara* from Ayvalık had modest activity on VRE (70%), however same species collected from Fethiye had strong activity both against MRSA (94%) and VRE (87%). While *Agelas oroides* from Fethiye had strong activity against MRSA (98%), VRE (96%), PV (95%), another *Agelas oroides* species collected from Kaş shows strong activity against VRE (91%) and modest activity against MRSA (66%). *Axinella verrucosa* showed strong activity against MRSA (100%) and VRE (100%). Some other sponge species, *Sarcotragus spinulosa* and *Dictyonella incisa* had modest activity against MRSA (61% and 82%, respectively). *Ircinia fasciculata* and *Chondrosia reniformis* collected from Kemer and Turgutreis demonstrated moderate

activity against MRSA (77%) and PV (51%), respectively.

Antimicrobial activity is observed only with 50 µg/mL filtered extracts of two sponge species. *Dysidea avara* from Fethiye had strong activity against VRE (96%) and *Ircinia fasciculata* from Kemer had modest activity (59%). A number of studies on secondary metabolites of the genera *Axinella*, *Dysidea* and *Ircinia* have been reported in the literature (14-21). According to the studies that have been conducted on these sponges, *Axinella* species contain terpenes, alkaloids and cyclopeptides, *Ircinia* genus possesses major metabolites such as linear furanoterpenes. All these compounds may contribute to antimicrobial activity. Avorol isolated from *Dysidea avara* has antimicrobial

Table 3. Species, localities and % inhibition values of filtrated extracts 1X (50 µg/mL)

ID	Taxonomy	Location	MRSA	VRE	CA	CA_S	PA*	PV*
1	<i>Ircinia oros</i>	Fethiye 1	6	17				
2	<i>Dictyonella incisa</i>	Fethiye 15	1	11				
3	<i>Petrocia ficiformis</i>	Ayvalık 1	7	18				
4	<i>Ircinia oros</i>	Ayvalık 2	5	2				
5	<i>Dysidea avara</i>	Ayvalık 3	9	2				
6	<i>Agelas oroides</i>	Ayvalık 4	0	7				
7	<i>Axinella verrucosa</i>	Ayvalık 5	8	7				
8	<i>Aplysina aerophoba</i>	Ayvalık 6	3	0				
9	<i>Chondrilla nucula</i>	Ayvalık 10	3	0				
10	<i>Aplidium elegans</i>	Ayvalık 11	0	0				
11	<i>Eucinella singularis</i>	Ayvalık 12	9	35				
12	<i>Agelas oroides</i>	Ayvalık 13	12	12				
13	<i>Axinella polypoides</i>	Danaadası (32m)	35	22				
14	<i>Petrocia ficiformis</i>	Danaadası	0	19				
15	<i>Petrocia ficiformis</i>	Fethiye 2	4	21				
16	<i>Agelas oroides</i>	Fethiye 3	4	0				
17	<i>Axinella damicornis</i>	Fethiye 4	0	0				
18	<i>Axinella verrucosa</i>	Fethiye 5	0	13				
19	<i>Dysidea avara</i>	Fethiye 7	4	96				
20	<i>Sarcotragus spinulosa</i>	Fethiye 10	15	0				
21	<i>Axinella polypoides</i>	Fethiye 11	15	36				
22	<i>Ircinia variabilis</i>	Fethiye 13	23	23				
23	<i>Petrocia ficiformis</i>	Fethiye 14	21	33				
24	<i>Petrocia ficiformis</i>	Fethiye 14	9	33				
25	<i>Dictyonella incisa</i>	Fethiye 15	0	13				
26	<i>Chondrilla nucula</i>	Güvercinlik	10	0				
27	<i>Dysidea avara</i>	İbrice 4	0	39				
28	<i>Agelas oroides</i>	Kaş 2	1	37				
29	<i>Ciocalyptra carbolloi</i>	Kaş 5	35	32				
30	<i>Ircinia fasciculata</i>	Kemer 3	59	5				
31	<i>Cliona viridis</i>	Sinekli 8	40	24				
32	<i>Agelas oroides</i>	Turgutreis 17	0	30				
33	<i>Chondrosia reniformis</i>	Turgutreis 23	27	11				

properties. Considering secondary metabolites of *Dictyonella incisa*, only two studies describing the compound has been published (22, 23).

Results indicate that antimicrobial activity varies among sponge species. One of the reasons for this variety is the collection localities of the sponges. Same sponge species can show strong, modest or weak activity depending on their localities. Besides, the amounts of extracts play an important role as well. Activity results depend on the concentration of the extracts.

CONCLUSION

Upto date there have been many papers on chemistry of the genera *Axinella* and *Ircinia* which consist a part of this study, in contrast to *Dictyonella incisa* that have not been intensively studied yet. Therefore further investigation needs to be performed for compound(s) purified from this sponge species in order to elucidate its (their) chemical structure(s) for antimicrobial activity. In conclusion, it is believed that a rich source of antimicrobial drug candidates could be obtained from the secondary metabolites of marine sponges.

ACKNOWLEDGMENTS

This work was supported by Ankara University, Coordination Unit of Scientific Research Projects Office (09B3336005). Author are thankful to Martin Lanteigne (Canada, Charlottetown, Prince Edward University, Faculty of Science, Department of Chemistry) for conducting the bioassays.

REFERENCES

1. Faulkner DJ, Marine Natural Products, Nat Prod Rep 19, 1-48, 2002.
2. Sipkema D, Franssen MC, Osinga R, Tramper J, Wijffels RH, Marine sponges as pharmacy, Mar Biotechnol 7 (3), 142-62, 2005.
3. Mayer AM, Glaser KB, Cuevas C, Jacobs RS, Kem W, Little RD, McIntosh JM, Newman DJ, Potts BC, Shuster DE, The odyssey of marine pharmaceuticals: a current pipeline perspective, Trends Pharmacol Sci 31 (6), 255-265, 2010.
4. Keyzers RA, Davies-Coleman MT, Anti-inflammatory metabolites from marine sponges, Chem Soc Rev 34, 355-365, 2005.
5. Dembitsky VM, Glorizova TA, Poroikov VV, Novel antitumor agents: marine sponge alkaloids, their synthetic analogs and derivatives, Med Chem 5, 319-336, 2005.
6. Abad MJ, Bedoya L, Bermejo P, Marine compounds and the antimicrobial activities, Formatex 1293-1306, 2011.
7. Laport MS, Santos DC, Muricy G, Marine sponges: potential sources of new antimicrobial drugs, Curr Pharm Biotechnol 10, 86-105, 2009.
8. Marinho PR, Muricy GS, Silva MFL, Giambiagi-deMarval M, Laport MS, Antibiotic-resistant bacteria inhibited by extracts and fractions from Brazilian marine sponges, Bras Farmacog 20, 267-275, 2010.
9. Cariello I, Zanetti L, Cuomo V, Vanzanella F, Antimicrobial activity of avarol, a sesquiterpenoid hydroquinone from the marine sponge, Comp Biochem Physiol B 71, 281-283, 1982.
10. Thakur NL, Anil AC, Antibacterial activity of the sponge *Irciniaramosa*: Importance of its surface-associated bacteria, J Chem Ecol 26, 57-71, 2000.
11. Touati I, Chaieb K, Bakhrouf A, Gaddour K, Screening of antimicrobial activity of marine sponge extracts collected from Tunisian coast, J Mycol Med 17, 183-187, 2007.
12. Torres YR, Berlink RGS, Nascimento GG, Fortier SC, Pessoa C, de Moraes MO, Antibacterial activity against resistant bacteria and cytotoxicity of four alkaloid toxins isolated from the marine sponge *Arenosclera brasiliensis*, Toxicon 40, 885-891, 2002.
13. Tsoukatou M, Hellio C, Vagias C, Harvala C, Roussis V, Chemical defense and antifouling activity of three Mediterranean sponges of the genus *Ircinia*, Z Naturforsch 57c, 161D171, 2002.
14. Overy PD, Berrue F, Correa H, Hanif N, Hay K, Lanteigne M, Mquilian K, Duffy S, Boland P, Jagannathan R, Carr GS, Vansteeland M, Kerr RG, Sea foam as a source of fungal inoculum for the isolation of biologically active natural products, Mycology 5(3), 1-15, 2014.
15. Qia SH, Wang Y, Zhanga S, Steroids and alkaloids from the South China sea sponge *Axinella* sp, J Asian Nat Prod Res 11(12), 1040-1044, 2009.
16. Wätjen W, Putz A, Chovolou Y, Kampkötter A, Totzke F, Kubbutat MH, Proksch P, Konuklugil B, Hexahepta- and nonaprenylhydroquinones isolated from marine sponges *Sarcotragus muscarum* and *Ircinia fasciculata* inhibit NF-kappa B signalling in H4IIE cells, J Pharm Pharmacol 61(7), 919-924, 2009.
17. Forenza S., Minale L, Riccio R. New bromo-pyrrole derivatives from the sponges *Agelas-Oroides* J Chemi Soc Chem Comm 18, 1129-1130, 1971.
18. Fathi-Afshar R, Allen TM, Krueger CA, Cook DA, Clanachan AS, Vriend R, Baer HP, Cass CE, Some pharmacological activities of novel adenine related compounds isolated from marine sponges *Agelas mauritiana*, Can J Physiol Pharm 67, 276-281, 1989.
19. Carletti I, Long C, Funel C, Amade P. Yardenone A, New cytotoxic triterpenes from the Indian Ocean sponge *Axinella* cf. *bidderi*, J Nat Prod 66, 25-29, 2003.
20. Bickmeyer U, Assmann M, Köck M, Schütt CA, Secondary metabolite, 4,5-dibromopyrrole-2-carboxylic acid, from marine sponges of the genus *Agelas* alters cellular calcium signals, Environ Toxicol Pharmacol 19, 423-427, 2005.

21. Bickmeyer U, Grube A, Klings KW, Köck M, Ageladine A, A pyrrole-imidazole alkaloid from marine sponges, is a pH sensitive membrane permeable dye, *Biochem Bioph Res Co* 373, 419-422, 2008.
22. Hertiani T, Edrada-Ebel R, Ortlepp S, Van Soest RW, de Voogd NJ, Wray V, Hentschel U, Kozytska S, Müller WE, Proksch P, From anti-fouling to biofilm inhibition: new cytotoxic secondary metabolites from two Indonesian *Agelas* sponges, *Bioorg Med Chem* 18 (3), 1297-1311, 2010.
23. Ciminieuo P, Fattorusso E, Magno S, Mangoni A, A novel conjugated ketosteroid from the marine sponge *Dictyonella incisa*, *J Nat Prod* 52 (6), 1331-1333, 1989.
24. Mansoor TA, Hong J, Lee CO, Bae SJ, Im KS, Jung JH. Cytotoxic sterol derivatives from a marine sponge *Homaxinella* sp. *J Nat Prod* 68 (3), 331-336, 2005.

Received: 25.12.2014

Accepted: 09.04.2015

