



Determination of the Phototoxicity Potential of Commercially Available Tattoo Inks Using the 3T3-neutral Red Uptake Phototoxicity Test

Piyasada Satılan Dövme Boyalarının 3T3-nötral Kırmızısı Alım Fototoksiste Testi ile Fototoksiste Potansiyelinin Belirlenmesi

Elif Gözde UTKU TÜRK, Ayşe Tarbin JANNUZZI, Buket ALPERTUNGA*

Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Toxicology, Istanbul, Turkey

ABSTRACT

Objectives: Tattooing is an ancient practice and its popularity has been increasing in the recent years. After tattooing, complications may occur related to compose tattoo inks. In this study, the phototoxicity potential of the blue, red and black colors of the most commonly used three different commercially-available tattoo ink brands have been examined by performing *in vitro* 3T3-neutral red uptake (NRU) phototoxicity test.

Materials and Methods: In the study, the phototoxicity of serial diluted concentrations of tattoo inks were evaluated with *in vitro* 3T3-NRU phototoxicity test method according to OECD guide 432. The data obtained from the NRU test result were uploaded to Phototox software (version 2.0) and the phototoxicity potentials of tattoo inks were determined *via* the calculation of the mean photo effect (MPE) and photo irritation factor (PIF) values.

Results: The red, black and blue colors of three different commercially available tattoo inks did not cause a cytotoxic activity on BALB/c 3T3 cells with 3T3-NRU test. The IC_{50} values could not be determined +ultraviolet (UV) and -UV conditions. PIF values could not be calculated and MPE values were <0.1 , which predicts the absence of phototoxic effect for all of the tested tattoo inks.

Conclusion: All tested inks were evaluated as non-phototoxic according to the results of MPE values calculated using Phototox software. However, test results should be verified by other phototoxicity test methods to obtain a comprehensive evaluation of phototoxic complications of different tattoo inks.

Key words: Phototoxicity, *in vitro* phototoxicity, 3T3-NRU phototoxicity test, tattoo ink, BALB/c 3T3 cells

ÖZ

Amaç: Dövme yapmak eski zamanlardan beri süregelen bir uygulamadır ve popüleritesi son yıllarda artmaktadır. Dövme yapılmasından sonra dövme boyaalarının bileşimine bağlı olarak çeşitli komplikasyonlar oluşabilir. Bu çalışmada, piyasada en çok kullanılan üç farklı dövme boyası markasının mavi, kırmızı ve siyah renklerinin fototoksiste potansiyeli *in vitro* 3T3-nötral kırmızı alım (NRU) fototoksiste testi yapılarak incelenmiştir.

Gereç ve Yöntemler: Çalışmada, dövme boyaalarının seri olarak seyreltilmiş konsantrasyonlarının fototoksitesi, OECD kılavuz 432'ye göre *in vitro* 3T3-NRU fototoksiste test yöntemi ile değerlendirilmiştir. NRU test sonucundan elde edilen veriler Phototox yazılımıyla (sürüm 2.0) değerlendirilmiştir. Dövme boyaalarının fototoksiste potansiyelleri medyan foto etki (MPE) ve foto iritasyon faktörü (PIF) değerlerinin hesaplanmasıyla belirlenmiştir.

Bulgular: Ticari olarak temin edilebilen üç farklı dövme boyasının kırmızı, siyah ve mavi renkleri, 3T3-NRU testi ile BALB/c 3T3 hücrelerinde sitotoksik aktiviteye neden olmamıştır. +Ultraviyole (UV) ve -UV koşullarında IC_{50} değerleri belirlenmemiştir. PIF değerleri hesaplanamamıştır ve MPE değerleri $<0,1$ olarak belirlenmiştir. Bu sonuçlar test edilen tüm dövme boyaalarının fototoksik etkilerinin olmadığını göstermektedir.

Sonuç: Phototox yazılımı ile hesaplanan MPE değerlerinin sonuçlarına göre test edilen tüm boyaaların fototoksik olmadığı sonucuna ulaşılmıştır. Bununla birlikte, farklı dövme boyaalarının fototoksik komplikasyonlarının kapsamlı bir değerlendirmesini yapabilmek için test sonuçlarının diğer fototoksiste test yöntemleriyle doğrulanması gerekmektedir.

Anahtar kelimeler: Fototoksiste, *in vitro* fototoksiste, 3T3-NRU fototoksiste testi, dövme boyası, BALB/c 3T3 hücreleri

*Correspondence: tunga@istanbul.edu.tr, Phone: +90 532 251 48 88, ORCID-ID: orcid.org/0000-0001-6043-7560

Received: 02.05.2021, Accepted: 18.06.2021

©Turk J Pharm Sci, Published by Galenos Publishing House.

INTRODUCTION

Tattooing has existed worldwide for many centuries and increases its popularity, especially among young people today. Tattooing is applied by injecting tattoo ink into the dermis layer that is 1.5-2 mm below the skin with the help of a needle.¹ Despite the increasing popularity of tattooing, the toxicity profile of tattoo inks and the potential risks of these inks remain unknown. Tattoo inks have different formulations. Generally, tattoo inks are prepared by suspending pigments in a solvent. Apart from pigments and solvents, tattoo inks contain additives such as binding agents, preservatives, thickeners, and antioxidants.² One of the most significant problems in the toxicological evaluation of tattoo inks is the lack of information about the ink compositions. Additionally, there is no regulation on their use in tattoo inks for organic/inorganic pigments, carbon black, and different chemicals used as coloring agents in tattoo inks.³ Dermatological severe complications may occur during tattooing and there is a serious increase in the number of patients who apply to the dermatology doctor due to skin conditions caused by a tattoo. Allergic skin reactions caused by tattoos are the most common skin problem. Especially, red tattoo ink can cause allergic skin reactions.⁴

Phototoxicity describes the toxic response of the skin after light exposure due to chemicals present on the skin following cutaneous or systemic application. To produce a phototoxic effect, a chromophore or a photosensitizing molecule must absorb photons. The absorbed photon causes reactive oxygen species (ROS) formation and because of increased ROS level, symptoms such as edema, burning, and pain occur in the skin.^{5,6} The European Union banned animal testing in cosmetic products in 2009 and their import and sale in 2013 to reduce the number of laboratory animals used and protect animals from unnecessary pain and injuries.^{7,8} Subsequently, an increasing number of countries worldwide including Turkey have adopted the ban on animal testing for cosmetics.⁹ Therefore, the need for validated relevant alternative *in vitro* test methods has increased to make the toxicologic evaluation of cosmetic products.¹⁰ *In vivo* experiments to determine acute phototoxicity are not allowed in Europe since 2000. Instead, the validated and regulatory accepted 3T3-neutral red uptake (NRU) phototoxicity test is primarily used as an *in vitro* alternative method.¹¹ The 3T3-NRU allows users to test many factors such as different test material concentrations, exposure times, and ultraviolet (UV) irradiation dose. Also, it has been determined to be reliable as results obtained with *in vivo* acute phototoxicity tests in animals and humans.^{12,13}

Therefore, the study aimed to evaluate the phototoxic potential of the black, blue, and red colors of three different commercially available tattoo ink brands that are widely used by *in vitro* 3T3-NRU according to the Organisation for Economic Cooperation and Development (OECD) 432 guideline.

MATERIALS AND METHODS

Cell culture

The BALB/c 3T3 fibroblasts were purchased from ATCC. The cells were cultured in Dulbecco's modified Eagle's medium

(DMEM) supplemented with 10% heat-inactivated fetal bovine serum (FBS) and 1x antibiotic-antimycotic solution (all from Gibco) in a humidified atmosphere at 37°C.

Test compounds

The black, red and blue colors of three commercially available tattoo inks were used for the tests. Black-triple black, red-lipstick red, blue-muter earth (Eternal Tattoo Supply, Brighton, MI, USA), black-true black, red-bright red, blue-mario's light blue (Intenze Products Inc. NJ, USA), black, red, blue (Tang Dragon Tattoo, China) was used. Chlorpromazine was used as the positive control (Eczacıbaşı, Turkey).

The absorption spectrum of the tattoo inks

Before starting the phototoxicity test, the absorption spectra of the tattoo inks were measured in the 250-700 nm range (Epic, Biotek). For this purpose, tattoo inks were dissolved in DMEM medium without phenol red at 1% final concentration, and this diluent was used as blank.

Photosensitivity of the BALB/c 3T3 fibroblasts

To determine the photosensitivity of the cells, the BALB/c 3T3 fibroblasts were seeded on clear 96-well plates at a seeding density of 1×10^4 in DMEM without phenol red containing 10% FBS and 1x antibiotic antimycotic solution. The day after, cells were irradiated with 0-2.5-5-10-15 joule/cm² of UVA radiation. The UVA sensitivity of the cells was evaluated with NRU cell viability assay after 24 h, as mentioned in detail below.

Neutral red uptake phototoxicity assay

The BALB/c 3T3 fibroblast NRU assay was carried out as described using the OECD 432 guideline with minor modifications.¹³ Extra washing steps due to the high coloring of tattoo inks were added to the assay. Briefly, the BALB/c 3T3 cells were seeded on clear 96-well plates at a seeding density of 1×10^4 in DMEM without phenol red with 10% FBS and 1x antibiotic antimycotic solution (assay medium). After culturing the cells in the plate for 24 h, wells were washed with 1x PBS. Briefly, eight logarithmic dilution series of tattoo inks starting from 200 µg/mL to 1 µg/mL with assay medium was prepared. The prepared concentrations of tattoo inks were added to wells and allowed to incubate for 1 h. 2 different sets of plates (-UV and +UV) were prepared for the assay. Then, +UV plates were irradiated with 5 joule/cm² of UVA radiation from a Philips PL-L UVA lamp in a home-designed and constructed wooden box. UVA radiation calculation was made with a UVA light meter (lutron UVA-365SD, Taiwan). While +UV plates were irradiating, the -UV plates were kept in the dark. After irradiation, the solutions were discarded, and wells were washed with 1x PBS. Then, the cells were allowed to incubate overnight with the assay medium. To determine the phototoxic effects of the tattoo inks, the medium was replaced with 50 µg/mL-neutral red in assay medium and incubated for 3 h in a humidified atmosphere at 37°C. Then, the medium was discarded, and the uptaken neutral red was dissolved with a mixture of acetic acid, water and ethanol (1:49:50). The absorbance was read at 540 nm with a plate reader. Then, the phototoxicity of the tattoo inks was evaluated with Phototox version 2.0 software (ZEBET

Germany) by calculating the photo irritation factor (PIF) and the mean photo effect (MPE). Since the tattoo ink dilutions were prepared with DMEM without phenol red with 10% FBS and 1x antibiotic antimycotic solution (assay medium) and there was not any different solvent effect on the cells, only assay medium containing wells were used as the negative control. Therefore, negative control results used in the negative control sections in the Phototox program and IC_{50} , PIF, MPE values were not calculated for negative controls. According to the OECD guideline, MPE <0.1 predicts the absence of a phototoxic effect. MPE >0.1 and 0.15 predict a probable phototoxic effect and MPE >0.15 predicts a phototoxic effect.¹³

Statistical analysis

Phototox version 2.0 software (ZEBET Germany) was used for concentration-response analysis. Results are presented as mean standard deviation of three independent experiments run in triplicate.

RESULTS

According to OECD guideline 432, a substance must show absorption in the UV/visible area to be photoreactive.¹³ For this reason, absorption spectra of each ink were taken before starting the 3T3-NRU phototoxicity test. Absorption peaks of inks were found to be between 410 and 420 nm in black inks, 610-640 nm in blue inks and 560-570 nm in red inks.

Photosensitivity study results showed that increasing doses of UVA had a phototoxic effect on the BALB/c 3T3 fibroblasts. The viability of the 5 joule/cm² of UVA irradiated cells was $95.7\% \pm 2.36$ compared with non-irradiated cells (0 joule/cm²) and $>80\%$ viability of the cells meets the quality criteria of OECD test guideline no: 432, (Figure 1).¹³ To check the accuracy of the 3T3-NRU phototoxicity test under our laboratory conditions, an experiment was performed with chlorpromazine, which was selected as a positive control, in line with OECD guidelines. According to the OECD Guideline 432, chlorpromazine should have a PIF value greater than 14.4, and the MPE value should be in the range of 0.33-0.63. The IC_{50} values for +UV and -UV




Figure 1. Photosensitivity of the BALB/c 3T3 fibroblast cells. ** $p < 0.01$, **** $p < 0.0001$

should be 0.1-2.0 $\mu\text{g/mL}$ and 7.0-90.0 $\mu\text{g/mL}$, respectively.¹³ The results of the 3T3-NRU phototoxicity test with positive control showed that the PIF value for chlorpromazine is 27.7 ± 4.2 , the MPE value is 0.50 ± 0.11 , the IC_{50} values are 1.7 ± 0.28 $\mu\text{g/mL}$ at (+UV) and 47 ± 6.9 $\mu\text{g/mL}$ at (-UV). These values were found to comply with the OECD guideline 432 limits.

Tattoo inks phototoxicity evaluation with 3T3-NRU phototoxicity test is shown in Table 1. Based on our results, the red, black and blue colors of three different commercially available tattoo inks did not exhibit phototoxic potential with the 3T3-NRU phototoxicity test. The IC_{50} values could not be determined +UV and -UV conditions since tattoo inks did not show cytotoxic activity on BALB/c 3T3 cells. Consequently, PIF values could not be calculated, and MPE values were <0.1 , which predicts the absence of phototoxic effect, (Table 1).

DISCUSSION

While tattooing has been practiced throughout the world for many centuries, the popularity of tattooing has increased significantly recently due to decorative reasons. Inflammatory, infectious and neoplastic complications may be seen after tattooing and a part of the beforementioned complications is related to allergy and hypersensitivity to tattoo inks.¹⁴ Additionally, sensitivity to the sun around the tattooed area is common and 20% of individuals suffer from tattoo-associated complications.⁶ Despite the increasing number of tattooed individuals, there are not sufficient toxicological and pharmacokinetic evaluations of the intradermal use of inks and colorants used for tattooing.³

Tattoo inks mainly contain pigments, dyes, water, solvent additives such as glycerin, alcohol, and ethylene glycol, preservatives, stabilizers, and pH regulators.¹⁵ The coloring agents that are used in the inks can vary depending on the brand and the color.² The tattoo ink manufacturers not must disclose the chemical composition and exact ingredients in their inks that cause potential uncertainty in the evaluation of the toxic effects of the pigments, solvents, and binders that are used in the tattoo inks.¹

The black tattoo inks mainly contain carbon. Besides, black tattoo inks may contain mutagenic and carcinogenic compounds such as carbon black, polyaromatic aromatic hydrocarbons (PAHs), and phenols. PAHs under UV irradiation can generate singlet oxygen, which can result in tattoo-associated complications.¹⁵ The blue inks may contain elements such as cobalt, aluminum, copper, and it has been reported that cobalt-containing ink compositions cause more skin reactions and irritation than copper-containing ink compositions. Red tattoo inks may contain azo pigments and elements such as cadmium and iron for coloring.¹⁶ Tattoo-related skin allergies are especially observed when red tattoo ink is used. Red tattoo-related skin allergies are supposed to be related to azo compounds.¹⁷ The azo compounds that are used as pigments, under high energetic radiation and heat can cause the production of the aromatic amines and amines considered allergic sensitizers.¹⁵ Also, often tattoo ink contains azo dyes with unknown compositions.^{11,18} The

Table 1. Phototoxic evaluation of the red, black and blue colors of three different commercially available tattoo inks with Phototox version 2.0 software (ZEBET Germany)

Substance	IC ₅₀		PIF ± SD	MPE ± SD	Evaluation
	-UV	+UV			
Clorpromazine (positive control)	47±6.9 µg/mL	1.7±0.28 µg/mL	27.7±4.2	0.50±0.11	Phototoxic
Eternal - black	-	-	1	-0.361±0.07	Non-phototoxic
Eternal - red	-	-	1	-0.351±0.02	Non-phototoxic
Eternal - blue	-	-	1	-0.229±0.01	Non-phototoxic
Intenze - black	-	-	1	-0.559±0.02	Non-phototoxic
Intenze - red	-	-	1	-0.159±0.01	Non-phototoxic
Intenze - blue	-	-	1	-0.126±0.01	Non-phototoxic
Tang dragon - black	-	-	1	-1.330±0.01	Non-phototoxic
Tang dragon - red	-	-	1	-1.155±0.04	Non-phototoxic
Tang dragon - blue	-	-	1	-0.535±0.03	Non-phototoxic

UV: Ultraviolet, PIF: Photo irritation factor, SD: Standard deviation, MPE: Mean photo effect

local metabolism of these unknown azo dyes into porphyrins may have photo-sensitizing effects on the skin.¹⁹

Phototoxicity describes the tissue reactions caused by light and it is the toxic response of the skin that develops due to the light exposure of a substance that is applied to the organism systemically or subcutaneously.¹³ The phototoxic reactions are characterized by skin irritation or exaggerated sunburn-like symptoms such as erythema, tenderness, pruritus and edema in patients.²⁰

The NRU test is based on the retention of the NRU dye in the viable cell's lysosomes, and the amount of the dye in the lysosomes is proportional to the cell viability.²¹ However the alterations in the lysosomal membranes may affect the NRU test results and when the NRU test is used to evaluate the toxicity of substances that specifically target lysosomes, results may indicate artificially high cytotoxicity. This can be considered the main limitation of the NRU test for cytotoxicity studies.^{22,23} The 3T3-NRU phototoxicity test is a validated and regulatory accepted test for predicting phototoxicity.¹³ Therefore, the study aimed to evaluate the phototoxic potential of the black, blue and red colors of the most widely used three different commercially available tattoo ink brands by *in vitro* NRU phototoxicity test according to OECD 432 guideline. It has been reported that red, blue, and black tattoos cause more sun-related complaints than other colors.¹⁹ A clear relationship between having tattoos and skin cancer development has not established today. There are case reports of the development of cancer types such as melanoma, basal cell carcinomas, squamous cell carcinomas, and keratoacanthomas in persons with tattoos. However, these cancer developments are most likely not only a result of having a tattoo but multifactorial.^{24,25}

Our results showed that the black, blue and red colors of three different commercially available brands were not phototoxic by *in vitro* NRU phototoxicity test while the chlorpromazine

(positive control) result was within the range recommended by OECD 432 guideline.

A recent study was conducted to evaluate the phototoxic effects of different tattoo pigments. In this study, cadmium sulfide, carbazole, cadmium selenide, mercury (II) sulfide, chromium oxide, and cobalt aluminate were examined using the 3T3-NRU phototoxicity test and 3D human reconstructed skin model. The results of this study showed that only carbazole and cadmium sulfide exhibited phototoxicity potential with the 3T3-NRU phototoxicity test, and this result could only be confirmed for carbazole with the 3D human skin model.²⁶ Similar to our study, the findings of this study do not address the phototoxicity potential of red [cadmium selenide and mercury (II) sulfide] and blue (cobalt aluminate) pigments.

Regensburger et al.²⁷ studied the 20 well-known PAHs content of 19 commercially available black tattoo inks. Many PAHs have been shown to be carcinogenic and have mutagenic activity. Furthermore, under UV irradiation they can generate singlet oxygen, which can contribute the tattoo-associated complications.^{18,28,29} In the study, they evaluated the phototoxicity of PAH extracts from black tattoo inks through mitochondrial activity in human keratinocytes and found that some extracts caused singlet oxygen generation with UVA irradiation which might be indicative of phototoxic reactions.²⁷

Another study is conducted to make *in vitro* and *in vivo* toxicological evaluations of the blue, green, red and black tattoo inks. According to the study, the red and green tattoo inks showed higher *in vitro* and *in vivo* toxicity due to containing azo compounds while black tattoo ink was found to be the safest.³⁰ Wamer and Yin³¹ evaluated cytotoxic and photocytotoxic activity of eighteen TiO₂ containing permanent makeup inks in human dermal fibroblasts as inhibition of colony formation. They did not determine cytotoxicity among inks but did observe

eight inks were phototoxic under UVA irradiation. It has been reported that the higher the PIF values obtained from the 3T3-NRU phototoxicity test results in the higher the probability of finding phototoxic *in vivo*. Thus, guideline thresholds should be reconsidered for the better translation of the results to *in vivo*. Additionally, the lack of a barrier system is a known limitation of the 3T3-NRU phototoxicity test for testing topical agents.³²

CONCLUSION

In conclusion, our study results do not indicate the phototoxic potential of examined black, blue, and red colors of three different commercially available brands. However, the 3T3-NRU phototoxicity test results should be verified by other phototoxicity test methods to evaluate the phototoxicity of tattoo inks correctly. Also, it should be considered that the data obtained because of the phototoxicity test do not provide information about other toxicological properties of tattoo inks. However, evidence from some studies suggests that tattoo ink compositions are variable, and some of this ink compositions can cause cytotoxic and phototoxic reactions. Thus, it is crucial to have better regulations on tattoo ink compositions to minimize their risk of complications.

Conflict of interest: No conflict of interest was declared by the authors. The authors are solely responsible for the content and writing of this paper.

REFERENCES

- Grant CA, Twigg PC, Baker R, Tobin DJ. Tattoo ink nanoparticles in skin tissue and fibroblasts. *Beilstein J Nanotechnol.* 2015;6:1183-1191.
- Dirks M. Making innovative tattoo ink products with improved safety: possible and impossible ingredients in practical usage. *Curr Probl Dermatol.* 2015;48:118-127.
- Laux P, Tralau T, Tentschert J, Blume A, Dahouk SA, Bäumler W, Bernstein E, Bocca B, Alimonti A, Colebrook H, de Cuyper C, Dähne L, Hauri U, Howard PC, Janssen P, Katz L, Klitzman B, Kluger N, Krutak L, Platzek T, Scott-Lang V, Serup J, Teubner W, Schreiber I, Wilkniß E, Luch A. A medical-toxicological view of tattooing. *Lancet.* 2016;387:395-402.
- Lerche CM, Heerfordt IM, Serup J, Poulsen T, Wulf HC. Red tattoos, ultraviolet radiation and skin cancer in mice. *Exp Dermatol.* 2017;26:1091-1096.
- Mang R, Stege H, Krutmann J. Mechanisms of phototoxic and photoallergic reactions in Contact Dermatitis (5th ed). Berlin; Springer. 2011;6:97-104.
- Serup J, Niels H, Linnet JT, Møhl B, Olsen O, Westh H. Tattoos—health, risks and culture. With introduction of the “seamless prevention” strategy. Copenhagen: The Council on Health and Disease Prevention. 2015.
- Union P. Regulation (EC) no 1223/2009 of the European Parliament and of the Council. *The OJEU.* 2009;342:59.
- EC (European Commission). Full EU ban on animal testing for cosmetics enters into force. Press release. 2013.
- Sreedhar D, Manjula N, Pise SA, Ligade V. Ban of cosmetic testing on animals: a brief overview. *Int J Cur Res Rev.* 2020;12:113-116.
- Vinken M. 3Rs toxicity testing and disease modeling projects in the European Horizon 2020 research and innovation program. *EXCLI J.* 2020;19:775-784.
- Liebsch M, Spielmann H, Pape W, Krul C, Deguercy A, Eskes C. UV-induced effects. *Altern Lab Anim.* 2005;(Suppl 1):131-146.
- Spielmann H, Balls M, Dupuis J, Pape WJ, Pechovitch G, de Silva O, Holzhütter HG, Clothier R, Desolle P, Gerberick F, Liebsch M, Lovell WW, Maurer T, Pfannenbecker U, Potthast JM, Csato M, Sladowski D, Steiling W, Brantom P. The International EU/COLIPA In Vitro Phototoxicity Validation Study: Results of Phase II (Blind Trial). Part 1: the 3T3 NRU phototoxicity test. *Toxicol In Vitro.* 1998;12:305-327.
- OECD. Test Guideline 432: *In Vitro* 3T3 NRU Phototoxicity Test. OECD Guidel Test Chem. 2019.
- Juhas E, English JC 3rd. Tattoo-associated complications. *J Pediatr Adolesc Gynecol.* 2013;26:125-129.
- Savitha AS. Composition of Tattoo. TATTOO-The Invaluable Compendium for Dermatologists. New Delhi; The Health Sciences; 2017;24.
- Forte G, Petrucci F, Cristaudo A, Bocca B. Market survey on toxic metals contained in tattoo inks. *Sci Total Environ.* 2009;407:5997-6002.
- Serup J. Seamless prevention of adverse events from tattooing: integrated strategy emphasising the customer-tattooist interaction. *Curr Probl Dermatol.* 2015;48:236-247.
- Shinohara MM. Complications of decorative tattoo. *Clin Dermatol.* 2016;34:287-292.
- Hutton Carlsen K, Serup J. Photosensitivity and photodynamic events in black, red and blue tattoos are common: A ‘Beach Study’. *J Eur Acad Dermatol Venereol.* 2014;28:231-237.
- Kim K, Park H, Lim KM. Phototoxicity: Its Mechanism and Animal Alternative Test Methods. *Toxicol Res.* 2015;31:97-104. Erratum in: *Toxicol Res.* 2015;31:321.
- Repetto G, del Peso A, Zurita JL. Neutral red uptake assay for the estimation of cell viability/cytotoxicity. *Nat Protoc.* 2008;3:1125-1131.
- Clothier R. The FRAME modified neutral red uptake cytotoxicity test. *Invitox Protocol (3a).* 1991.
- Cudazzo G, Smart DJ, McHugh D, Vanscheeuwijck P. Lysosomotropic-related limitations of the BALB/c 3T3 cell-based neutral red uptake assay and an alternative testing approach for assessing e-liquid cytotoxicity. *Toxicol In Vitro.* 2019;61:104647.
- Høgsberg T, Loeschner K, Löf D, Serup J. Tattoo inks in general usage contain nanoparticles. *Br J Dermatol.* 2011;165:1210-1218.
- Kluger N, Phan A, Debarbieux S, Balme B, Thomas L. Skin cancers arising in tattoos: coincidental or not? *Dermatology.* 2008;217:219-221.
- Kim SY, Seo S, Choi KH, Yun J. Evaluation of phototoxicity of tattoo pigments using the 3 T3 neutral red uptake phototoxicity test and a 3D human reconstructed skin model. *Toxicol In Vitro.* 2020;65:104813.
- Regensburger J, Lehner K, Maisch T, Vasold R, Santarelli F, Engel E, Gollmer A, König B, Landthaler M, Bäumler W. Tattoo inks contain polycyclic aromatic hydrocarbons that additionally generate deleterious singlet oxygen. *Exp Dermatol.* 2010;19:e275-e281.
- Bao L, Xu A, Tong L, Chen S, Zhu L, Zhao Y, Zhao G, Jiang E, Wang J, Wu L. Activated toxicity of diesel particulate extract by ultraviolet a radiation in mammalian cells: role of singlet oxygen. *Environ Health Perspect.* 2009;117:436-441.

29. Nisbet ICT, LaGoy PK. Toxic equivalency factors (TEFs) for polycyclic aromatic hydrocarbons (PAHs). *Regul Toxicol Pharmacol.* 1992;16:290-300.
30. Arl M, Nogueira DJ, Schweitzer Köerich J, Mottim Justino N, Schulz Vicentini D, Gerson Matias W. Tattoo inks: Characterization and *in vivo* and *in vitro* toxicological evaluation. *J Hazard Mater.* 2019;364:548-561.
31. Wamer WG, Yin JJ. Photocytotoxicity in human dermal fibroblasts elicited by permanent makeup inks containing titanium dioxide. *J Cosmet Sci.* 2011;62:535-547.
32. Ceridono M, Tellner P, Bauer D, Barroso J, Alépée N, Corvi R, De Smedt A, Fellows MD, Gibbs NK, Heisler E, Jacobs A, Jirova D, Jones D, Kandárová H, Kasper P, Akunda JK, Krul C, Learn D, Liebsch M, Lynch AM, Muster W, Nakamura K, Nash JF, Pfannenbecker U, Phillips G, Robles C, Rogiers V, Van De Water F, Liminga UW, Vohr HW, Watrelos O, Woods J, Zuang V, Kreysa J, Wilcox P. The 3T3 neutral red uptake phototoxicity test: practical experience and implications for phototoxicity testing--the report of an ECVAM-EFPIA workshop. *Regul Toxicol Pharmacol.* 2012;63:480-488.